Appendix D

Areas of Significant Conservation Value

The information contained in Appendix D has been obtained from Davidson, R J; Courtney, S P; Millar, I R; Brown, D A; Deans, N; Clerke, P R; Dix, J C; Lawless, P F; Mavor, S J; McRae, S M 1995: Ecologically important marine, freshwater, island and mainland areas from Cape Soucis to Ure River, Marlborough, New Zealand: Recommendations for Protection. Department of Conservation, Nelson/Marlborough Conservancy. Occasional Publication No 16, 286 p.

Coastal and Marine (Intertidal and Subtidal)

| Northern O're | | | | |
|---------------|--|---------------|------|---|
| Number | Site | Status | Page | Conservation Value |
| 1 /30 | Whites Bay, Cloudy Bay, East Coast | Regional | 68 | Highly modified but forest regenerating. Large variety of exposed east coast marine life, sandy coast, sand, tussock and spinifex (regionally unique community). Nationally unique regenerating coastal broad leaf. Cultural (Maori) dwarf false musk. Re-establishment of Pingao. |
| 1/30 | Rarangi, Cloudy Bay, East Coast | Regional | 68 | As above. |
| 1/31 | Wairau Lagoons, Cloudy Bay, East Coast | National | 70 | Largest biologically important estuary on east coast. 90 species avifauna - 27% of which are endangered vulnerable or rare. Royal spoonbill, wrybill, caspian tern, banded dotterel (vulnerable) black stilt (endangered). Waterfowl. 22 species of fish include two species flatfish, whitebait. Lagoon acts as nursery area. White bluffs dominant feature, boulder bank. |
| 1/31 | White Bluffs, Cloudy Bay, East Coast | National | 70 | As above. |
| 1/32 | Seaview, Clifford Bay, East Coast | National | 73 | Weeping broom (threatened). Scientific reserve (flora). |
| 1/34 | Cloudy Bay Hectors Dolphin Area, East Coast | National | 77 | Hectors dolphin (nationally large population). |
| 1/34 | Clifford Bay Hectors Dolphin Area, East Coast | National | 77 | As above. |
| 1/35 | Cape Campbell Kelp Beds and Shore Platforms, East Coast | Regional | 79 | <i>Macrocystis pyifera</i> (kelp) and, adjacent to Cape. Marine habitat. |
| 1/36 | Chancet Rocks, East Coast | International | 80 | Silicifed fossil sponges (genus Turonia). Chancet rocks. Pingao, banded dotterel (threatened), New Zealand fur seal haulout area. High degree of natural character. |
| 1/36 | Needles, East Coast | International | 80 | As above. |
| 1/36 | Mirza Creek, East Coast | Regional | 80 | As above. |